arXiv:1702.06421v1 [math.CA] 21 Feb 2017

DYNAMIC K-STRUVE SUMUDU SOLUTIONS FOR FRACTIONAL KINETIC EQUATIONS

K.S. NISAR, F.B.M. BELGACEM

Abstract. In this present study, we investigate solutions for fractional kinetic equations, involving k-Struve functions using Sumudu transform. The methodology and results can be considered and applied to various related fractional problems in mathematical physics.

1. Introduction

The Struve function $H_p(x)$ introduced by Hermann Struve in 1882, defined for $p \in \mathbb{C}$ by

$$H_p(x) := \sum_{k=0}^{\infty} \frac{(-1)^k}{\Gamma(k+3/2)\,\Gamma\left(k+p+\frac{3}{2}\right)} \left(\frac{x}{2}\right)^{2k+p+1}, \tag{1.1}$$

is the particular solutions of the non-homogeneous Bessel differential equations, given by,

$$x^2 y''(x) + x y'(x) + \left(x^2 - p^2\right) y(x) = \frac{4\left(\frac{x}{2}\right)^{p+1}}{\sqrt{\pi}\Gamma(p+1/2)}. \tag{1.2}$$

The homogeneous version of (1.2) have Bessel functions of the first kind, denoted as $J_p(x)$, for solutions, which are finite at $x = 0$, when p a positive fraction and all integers [6], while tend diverge for negative fractions,p. The Struve functions occur in certain areas of physics and applied mathematics, for example, in water-wave and surface-wave problems [1, 17], as well as in problems on unsteady aerodynamics [35]. The Struve functions are also important in particle quantum dynamical studies of spin decoherence [34] and nanotubes [38]. For more details about Struve functions, their generalizations and properties, the esteemed reader is invited to consider references, [7, 8, 18, 23, 29–33, 40]. Recently, Nisar et al. [21] introduced and studied various properties of k-Struve function $\mathtt{S}^{\mathtt{k}}_{\nu,c}$ defined by

$$\mathtt{S}^{\mathtt{k}}_{\nu,c}(x) := \sum_{r=0}^{\infty} \frac{(-c)^r}{\Gamma_{\mathtt{k}}(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2})\Gamma(r+\frac{3}{2})} \left(\frac{x}{2}\right)^{2r+\frac{\nu}{\mathtt{k}}+1}. \tag{1.3}$$

2010 *Mathematics Subject Classification.* 26A33, 44A20, 33E12.

Key words and phrases. Fractional kinetic equations, Sumudu transforms, k-Struve function, Fractional calculus .

The sumudu transform of $\mathtt{k}-$Struve function is given by

$$\begin{aligned} S\left[\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\right] &= \int_0^\infty e^{-t}\mathtt{S}^{\mathtt{k}}_{\nu,c}(ut)dt \\ &= \int_0^\infty e^{-t}\sum_{r=0}^{\infty}\frac{(-c)^r}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2}\right)\Gamma\left(r+\frac{3}{2}\right)}\left(\frac{ut}{2}\right)^{2r+\frac{\nu}{\mathtt{k}}+1}dt \\ &= \sum_{r=0}^{\infty}\frac{(-c)^r}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2}\right)\Gamma\left(r+\frac{3}{2}\right)}\int_0^\infty e^{-t}\left(\frac{ut}{2}\right)^{\frac{\nu}{\mathtt{k}}+2r}dt \\ &= \sum_{r=0}^{\infty}\frac{(-c)^r\Gamma\left(\frac{\nu}{\mathtt{k}}+2r+2\right)}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2}\right)\Gamma\left(r+\frac{3}{2}\right)}\left(\frac{u}{2}\right)^{\frac{\nu}{\mathtt{k}}+1+2r} \end{aligned}$$

Now, using

$$\Gamma_{\mathtt{k}}(\gamma) = \mathtt{k}^{\frac{\gamma}{\mathtt{k}}-1}\Gamma\left(\frac{\gamma}{\mathtt{k}}\right) \tag{1.4}$$

we have the following

$$S\left[\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\right] = \sum_{r=0}^{\infty}\frac{(-c)^r\Gamma\left(\frac{\nu}{\mathtt{k}}+2r+2\right)}{\mathtt{k}^{r+\frac{\nu}{\mathtt{k}}+\frac{1}{2}}\Gamma\left(r+\frac{\nu}{\mathtt{k}}+\frac{3}{2}\right)\Gamma\left(r+\frac{3}{2}\right)}\left(\frac{u}{2}\right)^{\frac{\nu}{\mathtt{k}}+1+2r}.$$

Denoting the left hand side by $G(u)$, we have

$$\begin{aligned} G(u) &= S\left[\mathtt{S}^{\mathtt{k}}_{\nu,c}(t);u\right] \\ &= \left(\frac{u}{2}\right)^{\frac{\nu}{\mathtt{k}}+1}k^{-\frac{1}{2}-\frac{\nu}{\mathtt{k}}}{}_2\Psi_2\left[\begin{array}{c}(\frac{\nu}{\mathtt{k}}+2,2),(1,1)\\(\frac{\nu}{\mathtt{k}}+\frac{3}{2},1),(\frac{3}{2},1)\end{array}\middle|-\frac{cu^2}{4\mathtt{k}}\right] \end{aligned} \tag{1.5}$$

and inverse Sumudu transform of $\mathtt{k}$-Struve function is given by

$$\begin{aligned} S^{-1}\left[\mathtt{S}^{\mathtt{k}}_{\nu,c}(x)\right] &= S^{-1}\left[\sum_{r=0}^{\infty}\frac{(-c)^r}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2}\right)\Gamma\left(r+\frac{3}{2}\right)}\left(\frac{u}{2}\right)^{\frac{\nu}{\mathtt{k}}+1+2r}\right] \\ &= \sum_{r=0}^{\infty}\frac{(-c)^r\left(\frac{1}{2}\right)^{\frac{\nu}{\mathtt{k}}+1+2r}}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\nu+\frac{3\mathtt{k}}{2}\right)\Gamma\left(r+\frac{3}{2}\right)}\frac{(t)^{\frac{\nu}{\mathtt{k}}+2r}}{\Gamma\left(\frac{\nu}{\mathtt{k}}+1+2r\right)} \end{aligned}$$

Using (1.4), we get

$$= \left(\frac{t}{2}\right)^{\frac{\nu}{\mathtt{k}}}k^{-\frac{1}{2}-\frac{\nu}{\mathtt{k}}}{}_1\Psi_3\left[\begin{array}{c}(1,1)\\(\frac{\nu}{\mathtt{k}}+\frac{3}{2},1),(\frac{3}{2},1),(\frac{\nu}{\mathtt{k}},2)\end{array}\middle|-\frac{ct^2}{4\mathtt{k}}\right] \tag{1.6}$$

In this paper, we consider (1.3) to obtain the solution of the fractional kinetic equations. Our methodology herein is based on Sumudu transform, [4, 5]. Fractional calculus is developed to large area of mathematics physics and other engineering applications [14, 14, 19, 22, 24–28, 41] because of its importance and efficiency. The fractional differential

equation between a chemical reaction or a production scheme (such as in birth-death processes) was established and treated by Haubold and Mathai [16], (also see [3, 9, 14]).

2. Solution of generalized fractional Kinetic equations for ĸ-Struve function

Let the arbitrary reaction described by a time-dependent quantity $N = (N_t)$. The rate of change $\frac{dN}{dt}$ to be a balance between the destruction rate $\mathfrak{d}$ and the production rate $\mathfrak{p}$ of N, that is, $\frac{dN}{dt} = -\mathfrak{d} + \mathfrak{p}$. Generally, destruction and production depend on the quantity N itself, that is,

$$\frac{dN}{dt} = -\mathfrak{d}\left(N_t\right) + \mathfrak{p}\left(N_t\right), \tag{2.1}$$

where N_t described by $N_t\left(t^*\right) = N\left(t - t^*\right), t^* > 0$. Another form of (2.1) is,

$$\frac{dN_i}{dt} = -c_i N_i\left(t\right), \tag{2.2}$$

with $N_i\left(t = 0\right) = N_0$, which is the number of density of species i at time $t = 0$ and $c_i > 0$. The solution of (2.2) is,

$$N_i\left(t\right) = N_0 e^{-c_i t}. \tag{2.3}$$

Integrating (2.2) gives,

$$N\left(t\right) - N_0 = -c.\ {}_0D_t^{-1} N\left(t\right), \tag{2.4}$$

where ${}_0D_t^{-1}$ is the special case of the Riemann-Liouville integral operator and c is a constant. The fractional form of (2.4) due to [16] is,

$$N\left(t\right) - N_0 = -c_0^{\upsilon} {}_0D_t^{-\upsilon} N\left(t\right), \tag{2.5}$$

where ${}_0D_t^{-\upsilon}$ defined as

$${}_0D_t^{-\upsilon} f\left(t\right) = \frac{1}{\Gamma\left(\upsilon\right)} \int_0^t \left(t - s\right)^{\upsilon - 1} f\left(s\right) ds, \Re\left(\upsilon\right) > 0. \tag{2.6}$$

Suppose that $f(t)$ is a real or complex valued function of the (time) variable $t > 0$ and s is a real or complex parameter. The Laplace transform of $f(t)$ is defined by

$$F\left(p\right) = L\left[f(t) : p\right] = \int_0^{\infty} e^{-pt} f\left(t\right) dt, \quad \Re\left(p\right) > 0 \tag{2.7}$$

The Mittag-Leffler functions $E_\alpha\left(z\right)$ (see [20]) and $E_{\alpha,\beta}\left(x\right)$ [39] is defined respectively as

$$E_\alpha\left(z\right) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma\left(\alpha n + 1\right)} \quad \left(z, \alpha \in \mathbb{C}; |z| < 0, \Re\left(\alpha\right) > 0\right). \tag{2.8}$$

$$E_{\alpha,\beta}(x)=\sum_{n=0}^{\infty}\frac{x^n}{\Gamma(\alpha n+\beta)}. \tag{2.9}$$

Theorem 1. *If $d>0,\nu>0,\mu,c,t\in\mathbb{C}$ and $l>-\frac{3}{2}\mathtt{k}$ then the solution of generalized fractional kinetic equation*

$$N(t)=N_0\ \mathtt{S}^{\mathtt{k}}_{\mu,c}(d^\nu t^\nu)-d^\nu\ {}_0D_t^{-\nu}N(t), \tag{2.10}$$

is given by the following formula

$$N(t)=N_0\sum_{r=0}^{\infty}\frac{(-c)^r\,\Gamma\left[\nu\left(2r+\frac{\mu}{\mathtt{k}}+1\right)+1\right]}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)}\frac{1}{t}\left(\frac{d^\nu t^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1}E_{\nu,\nu(2r+\frac{\mu}{\mathtt{k}})+1}(-d^\nu t^\nu). \tag{2.11}$$

where $E_{\nu,\nu(2r+\frac{\mu}{\mathtt{k}})+1}(-d^\nu t^\nu)$ is given in (2.9)

Proof. The Sumudu transform of Riemann-Lioville fractional integral operators is given by

$$S\left\{{}_0D_t^{-\nu}f(t);u\right\}=u^\nu G(u), \tag{2.12}$$

where $G(u)$ is defined in (1.5). Now applying Sumudu transform both sides of (2.10) and applying the definition of $\mathtt{k}$-Struve function given in (1.3), we have

$$\begin{aligned}N^*(u)&=S\left[N(t);u\right]\\&=N_0S\left[\mathtt{S}^{\mathtt{k}}_{\mu,c}(d^\nu t^\nu);u\right]-d^\nu S\left[{}_0D_t^{-\nu}N(t);u\right]\\&=N_0\left[\int_0^\infty e^{-pt}\sum_{r=0}^{\infty}\frac{(-c)^r}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)}\left(\frac{d^\nu(ut)^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1}dt\right]\\&-d^\nu u^\nu N^*(u),\end{aligned}$$

where

$$S\left\{t^{\mu-1}\right\}=u^{\mu-1}\Gamma(\mu). \tag{2.13}$$

By rearranging terms we get,

$$\begin{aligned}&N^*(u)+d^\nu u^\nu N^*(u)\\&=N_0\sum_{r=0}^{\infty}\frac{(-c)^r}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)}\left(\frac{d^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1}\\&\times\int_0^\infty e^{-t}(ut)^{\nu(2r+\frac{\mu}{\mathtt{k}}+1)}\,dt\end{aligned}$$

$$= N_0 \sum_{r=0}^{\infty} \frac{(-c)^r \Gamma[\nu(2r+\frac{\mu}{k}+1)+1]}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)} \left(\frac{u^\nu d^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1}$$

Therefore

$$N^*(u) = N_0 \sum_{r=0}^{\infty} \frac{(-c)^r \Gamma[\nu(2r+\frac{\mu}{k}+1)+1]}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)} \left(\frac{d^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1} \times \left\{ u^{\nu(2r+\frac{\mu}{\mathtt{k}}+1)} \sum_{n=0}^{\infty} [-(du)^\nu]^n \right\} \tag{2.14}$$

Taking inverse Sumudu transform of (2.14), and by using

$$S^{-1}\{u^\nu; t\} = \frac{t^{\nu-1}}{\Gamma(\nu)}, \Re(\nu) > 0, \tag{2.15}$$

we have

$$S^{-1}\{N^*(u)\} = N_0 \sum_{r=0}^{\infty} \frac{(-c)^r \Gamma[\nu(2r+\frac{\mu}{k}+1)+1]}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)} \left(\frac{d^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1} \times S^{-1}\left\{ \sum_{n=0}^{\infty} (-1)^n (d)^{\nu n} u^{\nu(2r+\frac{\mu}{\mathtt{k}}+n+1)} \right\}$$

which gives,

$$\begin{aligned} N(t) &= N_0 \sum_{r=0}^{\infty} \frac{(-c)^r \Gamma[\nu(2r+\frac{\mu}{\mathtt{k}}+1)+1]}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)} \left(\frac{d^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1} \\ &\times \left\{ \sum_{n=0}^{\infty} (-1)^n (d)^{\nu n} \frac{t^{\nu\left(2r+\frac{\mu}{\mathtt{k}}+n+1\right)-1}}{\Gamma\left[\nu\left(2r+\frac{\mu}{\mathtt{k}}+n+1\right)\right]} \right\} \\ &= N_0 \sum_{r=0}^{\infty} \frac{(-c)^r \Gamma[\nu(2r+\frac{\mu}{\mathtt{k}}+1)+1]}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)} \frac{1}{t} \left(\frac{d^\nu t^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1} \\ &\times \left\{ \sum_{n=0}^{\infty} (-1)^n (d)^{\nu n} \frac{t^\nu}{\Gamma\left[\nu\left(2r+\frac{\mu}{\mathtt{k}}+n+1\right)\right]} \right\} \\ &= N_0 \sum_{r=0}^{\infty} \frac{(-c)^r \Gamma\left[\nu\left(2r+\frac{\mu}{\mathtt{k}}+1\right)+1\right]}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)} \frac{1}{t} \left(\frac{d^\nu t^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1} \\ &\times E_{\nu,\nu(2r+\frac{\mu}{\mathtt{k}})+1}(-d^\nu t^\nu) \end{aligned}$$

which is the desired result. □

Corollary 2.1. *If we put* $\mathtt{k} = 1$ *in* (2.11) *then we get the solution of fractional kinetic equation involving classical Struve function as:*

If $d>0,\nu>0,\mu,c,t\in\mathbb{C}$ *and* $l>-\frac{3}{2}$ *then the solution of generalized fractional kinetic equation*

$$N(t)=N_0\ \mathtt{S}^1_{\mu,c}\left(d^\nu t^\nu\right)-d^\nu\ {}_0D_t^{-\nu}N(t)\,, \tag{2.16}$$

is given by the following formula

$$N(t)=N_0\sum_{r=0}^{\infty}\frac{(-c)^r\,\Gamma\left[\nu\left(2r+\mu+1\right)+1\right]}{\Gamma\left(r+\mu+\frac{3}{2}\right)\Gamma\left(r+\frac{3}{2}\right)}\frac{1}{t}\left(\frac{d^\nu t^\nu}{2}\right)^{2r+\mu+1}E_{\nu,\nu(2r+\mu)+1}\left(-d^\nu t^\nu\right). \tag{2.17}$$

Theorem 2. *If* $\mathfrak{a}>0,d>0,\nu>0,c,\mu,t\in\mathbb{C},\mathfrak{a}\neq d$ *and* $\mu>-\frac{3}{2}\mathtt{k}$*, then the equation*

$$N(t)=N_0\ \mathtt{S}^{\mathtt{k}}_{\mu,c}\left(\mathfrak{a}^\nu t^\nu\right)-d^\nu\ {}_0D_t^{-\nu}N(t)\,, \tag{2.18}$$

is given by the following formula

$$\begin{aligned}N(t)=N_0\sum_{r=0}^{\infty}&\frac{(-c)^r\,\Gamma\left[\nu\left(2r+\frac{\mu}{\mathtt{k}}+1\right)+1\right]}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)}\frac{1}{t}\left(\frac{d^\nu t^\nu}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1}\\&\times E_{\nu,\nu(2r+\frac{\mu}{\mathtt{k}})+1}\left(-\mathfrak{a}^\nu t^\nu\right).\end{aligned} \tag{2.19}$$

where $E_{\nu,\nu(2r+\frac{\mu}{\mathtt{k}})+1}(.)$ *is given in* (2.9)

Proof. Theorem 2 can be proved in parallel with the proof of Theorem 1. So the details of proofs are omitted. □

Corollary 2.2. *By putting* $\mathtt{k}=1$ *in Theorem 2, we get the solution of fractional kinetic equation involving classical Struve function: If* $\mathfrak{a}>0,d>0,\nu>0,c,\mu,t\in\mathbb{C},\mathfrak{a}\neq d$ *and* $\mu>-\frac{3}{2}$*, then the equation*

$$N(t)=N_0\ \mathtt{S}^1_{\mu,c}\left(\mathfrak{a}^\nu t^\nu\right)-d^\nu\ {}_0D_t^{-\nu}N(t)\,, \tag{2.20}$$

is given by the following formula

$$\begin{aligned}N(t)=N_0\sum_{r=0}^{\infty}&\frac{(-c)^r\,\Gamma\left[\nu\left(2r+\mu+1\right)+1\right]}{\Gamma\left(r+\mu+\frac{3}{2}\right)\Gamma\left(r+\frac{3}{2}\right)}\frac{1}{t}\left(\frac{d^\nu t^\nu}{2}\right)^{2r+\mu+1}\\&\times E_{\nu,\nu(2r+\mu)+1}\left(-\mathfrak{a}^\nu t^\nu\right).\end{aligned} \tag{2.21}$$

Theorem 3. *If* $d>0,\nu>0,c,\mu,t\in\mathbb{C}$ *and* $\mu>-\frac{3}{2}\mathtt{k}$*, then the solution of the following equation*

$$N(t)=N_0\ \mathtt{S}^{\mathtt{k}}_{\mu,c}\left(t^\nu\right)-d_0^\nu D_t^{-\nu}N(t)\,, \tag{2.22}$$

is given by the following formula

$$\begin{aligned}N(t)=N_0\sum_{r=0}^{\infty}&\frac{(-c)^r\,\Gamma\left[\nu\left(2r+\frac{\mu}{\mathtt{k}}+1\right)+1\right]}{\Gamma_{\mathtt{k}}\left(r\mathtt{k}+\mu+\frac{3}{2}\mathtt{k}\right)\Gamma\left(r+\frac{3}{2}\right)}\frac{1}{t}\left(\frac{t}{2}\right)^{2r+\frac{\mu}{\mathtt{k}}+1}\\&\times E_{\nu,\nu(2r+\frac{\mu}{\mathtt{k}})+1}\left(-d^\nu t^\nu\right).\end{aligned} \tag{2.23}$$

where $E_{\nu,\nu(2r+\frac{\mu}{\mathtt{k}})+1}(.)$ *is given in* (2.9)

Corollary 2.3. *If we set* $\mathtt{k}=1$ *then* (2.23) *reduced as follows:*
If $d>0,\nu>0,c,\mu,t\in\mathbb{C}$ *and* $\mu>-\frac{3}{2}$*, then the solution of the following equation*

$$N(t)=N_0\ \mathtt{S}^1_{\mu,c}(t^\nu)-d_0^\nu D_t^{-\nu}N(t)\,, \tag{2.24}$$

is given by the following formula

$$N(t)=N_0\sum_{r=0}^{\infty}\frac{(-c)^r\,\Gamma\left[\nu\left(2r+\mu+1\right)+1\right]}{\Gamma\left(r+\mu+\frac{3}{2}\right)\Gamma\left(r+\frac{3}{2}\right)}\frac{1}{t}\left(\frac{t}{2}\right)^{2r+\mu+1}$$
$$\times E_{\nu,\nu(2r+\mu)+1}\left(-d^\nu t^\nu\right). \tag{2.25}$$

3. Graphical interpretation

In this section we plot the graphs of our solutions of the fractional kinetic equation, which is established in (2.11). In each graph, we gave three solutions of the results on the basis of assigning different values to the parameters.In figures 1, we take $\mathtt{k}=1$ and $\nu=0.5,0.7,0.9,1,1.5$. Similarly figures 2-3 are plotted respectively by taking $\mathtt{k}=2,3$. Figures 4-6 are plotted by considering the solution given in (2.23) by taking $\nu=0.5,0.7,0.9,1,1.5$ and $\mathtt{k}=1,2,3$. Other than ν and $\mathtt{k}$ all other parameters are fixed by 1. It is clear from these figures that $N_t>0$ for $t>0$ and $N_t>0$ is monotonic increasing function for $t\in(0,\infty)$. In this study, we choose first 50 terms of Mittag-Leffler function and first 50 terms of our solutions to plot the graphs. $N_t=0$, when $t>0$ and $N_t\to\infty$ when $t\to1$ for all values of the parameters.

References

[1] A. R. Ahmadi and S. E. Widnall, Unsteady lifting-line theory as a singular-perturbation problem. J. Fluid Mech. 153 (1985), 59-81.

[2] Á. Baricz, Generalized Bessel functions of the first kind, Lecture Notes in Mathematics,1994, Springer, Berlin, 2010.

[3] F.B.M. Belgacem, A.A. Karaballi, S.L Kalla, Analytical Investigations of the Sumudu Transform and Applications to Integral Production Equations, Journal of Mathematical Problems in Engineering, No. 3, (2003), 103-118.

[4] F.B.M. Belgacem, A.A. Karaballi, Sumudu Transform Fundamental Properties Investigations and Applications, Journal of Applied Mathematics and Stochastic Analysis, Vol. (2006), Article ID 91083.

[5] F.B.M. Belgacem, Introducing and Analyzing Deeper Sumudu Properties, Nonlinear Studies, Vol. 13, No.1, (2006), 23-42.

[6] F.B.M. Belgacem, Applications with the Sumudu Transform to Bessel Functions and Equations, Applied Math. Sciences, Vol. 4, No. 74, (2010), 3665-3686.

[7] K. N. Bhowmick, Some relations between a generalized Struve's function and hypergeometric functions, Vijnana Parishad Anusandhan Patrika 5 (1962), 93-99.

[8] K. N. Bhowmick, A generalized Struve's function and its recurrence formula, Vijnana Parishad Anusandhan Patrika 6 (1963), 1-11.
[9] V.B.L. Chaurasia and S.C. Pandey, On the new computable solution of the generalized fractional kinetic equations involving the generalized function for the fractional calculus and related functions, Astrophys. Space Sci., 317 (2008), 213-219.
[10] A. Chouhan, S.D. Purohit and S. Saraswat, An alternative method for solving generalized differential equations of fractional order, Kragujevac J. Math., 37(2) (2013), 299-306.
[11] A. Chouhan, S. Sarswat, On solution of generalized Kinetic equation of fractional order, Int. Jr. of Mathematical Sciences and Applications, 2 (2)(2012), 813-818.
[12] A. Erdélyi, W. Magnus, F. Oberhettinger, and F.G. Tricomi, Tables of Integral Transforms, Vol. 1, McGraw-Hill, New York-Toronto-London, 1954.
[13] L. Galué, A generalized Bessel function, Integral Transforms Spec. Funct. 14 (2003), no. 5, 395-401.
[14] V.G. Gupta, B. Sharma and F. B. M. Belgacem, On the solutions of generalized fractional kinetic equations, Applied Mathematical Sciences, 5(19) (2011), 899-910.
[15] A. Gupta and C.L. Parihar, On solutions of generalized kinetic equations of fractional order, Bol. Soc. Paran. Mat., 32(1) (2014), 181-189.
[16] H.J. Haubold and A.M. Mathai, The fractional kinetic equation and thermonuclear functions, Astrophys. Space Sci., 327 (2000), 53-63.
[17] M. H. Hirata, Flow near the bow of a steadily turning ship, J. Fluid Mech. 71 (1975), no. 2, 283-291.
[18] B. N. Kanth, Integrals involving generalized Struve's function, Nepali Math. Sci. Rep. 6 (1981), no. 1-2, 61-64.
[19] Q.K. Katatbeh, F.B.M. Belgacem, Applications of the Sumudu Transform to Fractional Differential Equations, Nonlinear Studies, Vol. 18, No.1, (2011), 99-112.
[20] G.M. Mittag-Leffler, Sur la representation analytiqie d'une fonction monogene cinquieme note, Acta Math., 29 (1905), 101-181.
[21] K.S Nisar, S.R Mondal and J. Choi, Certain inequalities involving the k-Struve function, submitted, 2016
[22] K.S Nisar, S.D Purohit and S.R Mondal, Generalized Fractional Kinetic equations involving generalized Struve function of the first Kind, Journal of King Saud University-Science (2015), http://dx.doi.org/10.1016/j.jksus.2015.08.005.
[23] H. Orhan and N. Yagmur, Geometric Properties of Generalized Struve Functions, An. Ştiinţ . Univ. Al. I. Cuza Ia si. Mat. (N.S.), DOI: 10.2478/aicu-2014-0007, May 2014.
[24] A. Saichev and M. Zaslavsky, Fractional kinetic equations: solutions and applications, Chaos, 7(1997), 753-764.
[25] R.K. Saxena, A.M. Mathai, and H.J. Haubold, On Fractional kinetic equations, Astrophys. Space Sci., 282 (2002), 281-287.
[26] R.K. Saxena, A.M. Mathai, and H.J. Haubold, On generalized fractional kinetic equations, Physica A 344 (2004), 657-664.
[27] R.K. Saxena, A.M. Mathai and H.J. Haubold, Solution of generalized fractional reaction-diffusion equations, Astrophys. Space Sci., 305 (2006), 305-313.
[28] R.K. Saxena and S.L. Kalla, On the solutions of certain fractional kinetic equations, Appl. Math. Comput., 199 (2008), 504-511.

[29] R. P. Singh, Generalized Struve's function and its recurrence relations, Ranchi Univ. Math. J. 5 (1974), 67-75.
[30] R. P. Singh, Generalized Struve's function and its recurrence equation, Vijnana Parishad Anusandhan Patrika 28 (1985), no. 3, 287-292.
[31] R. P. Singh, Some integral representation of generalized Struve's function, Math. Ed. (Siwan) 22 (1988), no. 3, 91-94.
[32] R. P. Singh, On definite integrals involving generalized Struve's function, Math. Ed. (Siwan) 22 (1988), no. 2, 62-66.
[33] R. P. Singh, Infinite integrals involving generalized Struve function, Math. Ed. (Siwan) 23 (1989), no. 1, 30-36.
[34] J. Shao, P. Hanggi, Decoherent dynamics of a two-level system coupled to a sea of spins. Phys. Rev. Lett. 81 (1998), no. 26, 5710-5713.
[35] D. C. Shaw, Perturbational results for diraction of water-waves by nearly-vertical barriers. IMA, J. Appl. Math. 34 (1985), no. 1, 99-117.
[36] H.M. Srivastava and R.K. Saxena, Operators of fractional integration and their applications, Appl. Math. Comput., 118 (2001), 1-52.
[37] M.R. Spiegel, Theory and Problems of Laplace Transforms, Schaums Outline Series, McGraw-Hill, New York, 1965.
[38] T. G. Pedersen, Variational approach to excitons in carbon nanotubes. Phys. Rev. B 67 (2003), no. 7, (0734011)-(0734014).
[39] A. Wiman, Uber den fundamental satz in der theorie der funktionen $E_{\alpha}(z)$, Acta Math. 29 (1905), 191-201.
[40] N. Yagmur and H. Orhan, Starlikeness and convexity of generalized Struve functions, Abstr. Appl. Anal. 2013, Art. ID 954513, 6 pp.
[41] G.M. Zaslavsky, Fractional kinetic equation for Hamiltonian chaos, Physica D, 76(1994), 110-122.

K. S. Nisar : Department of Mathematics, College of Arts and Science at Wadi Al-dawaser, Prince Sattam bin Abdulaziz University, Riyadh region 11991, Saudi Arabia
E-mail address: ksnisar1@gmail.com, n.sooppy@psau.edu.sa

F.B.M. Belgacem: Department of Mathematics, Faculty of Basic Education, PAAET, Al-Ardhiya, Kuwait
E-mail address: fbmbelgacem@gmail.com